

School Asthma Action Plan

Name	Birth Date
Emergency Contact	Phone Cell
Triggers	<input type="checkbox"/> Mold/Pollens <input type="checkbox"/> Animals <input type="checkbox"/> Colds <input type="checkbox"/> Dust <input type="checkbox"/> Exercise <input type="checkbox"/> Smoke <input type="checkbox"/> Weather <input type="checkbox"/> Fragrance

Green Zone: Doing Well

- Breathing is good • No cough or wheeze • Can work and play • Sleeps all night
- No early warning signs • Peak Flow Meter if used: 80-100% of personal best _____

School Action: Follow actions in marked boxes below for exercise induced asthma

Medication Before Exercise
 Medication Before Recess
 Use routinely every _____ hours
Medication with spacer: Albuterol Ventolin Proventil Xopenex
Medication without spacer: Maxair Autohaler
Dose: _____ puffs
 When: 10-15 minutes before listed activity
 Start Date: School Year
 Stop Date: School Year

Yellow Zone: Getting Worse
(mild trouble breathing)

- Cough, wheeze, chest tight • Problems working/ playing
- Early warning signs • Shortness of breath
- Peak Flow Meter if used: 50 to 80% of personal best _____

School Actions: Follow actions in marked boxes below

Take Quick-Relief Medication	How Much (Dose)	When	Start Date	Stop Date
MDI with Spacer: <input type="checkbox"/> Albuterol <input type="checkbox"/> Ventolin <input type="checkbox"/> Proventil <input type="checkbox"/> Xopenex Without spacer: <input type="checkbox"/> Maxair Autohaler	_____ puffs	Student report of symptoms	School Year	School Year
Nebulizer: <input type="checkbox"/> Albuterol <input type="checkbox"/> Ventolin <input type="checkbox"/> Proventil <input type="checkbox"/> Xopenex	_____ Unit Dose	Student report of symptoms	School Year	School Year

- If symptoms improve after 10-15 minutes: Return to normal activity
- If symptoms do not improve after 10-15 minutes: Give quick relief medication again and call parents
- If symptoms improve after the second 10-15 minutes: Return to normal activity and call parents
- If symptoms do not improve after the medication is repeated: Call EMS (911), School RN and parents
- If symptoms get worse at anytime: Call EMS (911), School RN and Parents
- Report frequent use of quick relief medications (twice a day for 3 days, not for exercise) to the School RN and Parents

Red Zone: Medical Alert
(severe trouble breathing)

- Cannot stop coughing • Breathing fast • Flaring nostrils • Medication not helping
- Getting worse, instead of better • Trouble walking or talking from shortness of breath
- The skin between the ribs and above the collarbone pulls in or retracts
- Lips or fingernails are blue

School Actions:

1. Call EMS (911) IMMEDIATELY
2. GIVE QUICK-RELIEVER MEDICATION AND CONTINUE EVERY 15 MINUTES UNTIL EMS (911) ARRIVES
3. Call School RN and Parents

Take Quick-Relief Medications	How Much (Dose)	When	Start Date	Stop Date
MDI with Spacer: <input type="checkbox"/> Albuterol <input type="checkbox"/> Ventolin <input type="checkbox"/> Proventil <input type="checkbox"/> Xopenex Without spacer: <input type="checkbox"/> Maxair Autohaler	_____ puffs	Student report of or observation of symptoms.	School Year	School Year
Nebulizer: <input type="checkbox"/> Albuterol <input type="checkbox"/> Ventolin <input type="checkbox"/> Proventil <input type="checkbox"/> Xopenex	_____ Unit Dose	Student report of or observation of symptoms	School Year	School Year

Health Care Provider Name: _____
 Phone: _____
 FAX#: _____

Health Care Provider Sign: _____
 Date: _____

Student Name: _____ Birth Date: _____

School Asthma Action Plan (page two)

Metered Dose Inhaler (MDI) Instructions

1. Store at room temperature.
2. Shake the MDI for 5 seconds before each use.
3. Prime the MDI before the first use or when not used every day. Follow the product's patient information sheet for MDI specific priming instructions. Priming usually involves pressing down on the medication canister to discard into the air one or more puffs of medication. Discarding puffs makes sure the next puff inhaled contains the labeled amount of medication.
3. Keep track of metered inhalation puffs used. Subtract the number used from the number of metered inhalation puffs available listed on the label. The number of metered inhaled puffs available is listed on the medication canister or on the box. There are usually 200 puffs in an MDI.
4. Ask family for a new MDI when all labeled metered inhalation puffs are used.

MDI and Aerosol Solution Potential Adverse Reaction: Headache, shakiness, fast heart rate, nausea. Call parent with 1) student report of symptoms that interfere with schoolwork or activity 2) increase in side effects 3) frequent usage (2 times a day for 3 days).

We have instructed the patient and family in the proper use of the quick-relief medications. It is my professional opinion that the student:

_____ should be allowed to carry and self administer the inhaled medication.

_____ should **not** carry and self administer the inhaled medication. The medication should be stored and administered by designated school personnel.

Provider Signature Date

Section II To Be Completed by Parent

I give permission for my child to receive medication at school in keeping with Section I above according to the school district policy and as instructed by the physician and agree to 1) Assume responsibility for safe delivery of the medication in its original container to the school, 2) Have a new form completed by the doctor if medication or dosage is changed, 3) Notify the school of changes in health care provider. I release from liability, and in addition agree to indemnify, all school employees, the Board of Education and Akron Children's Hospital School Health Services for damages or injury resulting from the use, misuse or nonuse of such medication except as such Board, School Health Services or its employees are grossly negligent or engage in wanton or reckless misconduct. I further agree to submit a revised statement signed by the physician who has prescribed the medication described in Section I in the event that I become aware that any of the above information has changed. I have read and understand the policy of the School Board for administration of medication and affirm that this request entails special circumstances justifying an exception from the usual administration of medication at school-by-school personnel.

Parent/Guardian Signature: _____ Date: _____

Daytime Phone: _____

THIS FORM EXPIRES AT THE END OF THE SCHOOL YEAR